

# Venaseal<sup>™</sup> Post Procedure Instructions

### What to expect

- Recovery from Venaseal<sup>™</sup> closure procedure is usually trouble free. Typical symptoms resolve in 2-3 days and include:
  - Mild soreness at the vein access site
  - Mild discomfort, bruising or swelling along the treated vein
  - Redness over the treated area
  - Firm or cord-like area over the treated vein

## **Activity post-procedure**

- Routine activities such as walking, grocery shopping, stairs and light exercise (treadmill, cycling, aerobics, yoga, etc..) can be resumed immediately following the procedure
  - You should walk a minimum of 30 minutes daily (2-3 times 15 minutes each) for the first week following the procedure
- Avoid strenuous activity or heavy lifting (greater than 10-15lbs) for 7 days following procedure
- Avoid soaking in a hot tub, hot bath, Jacuzzi etc.. for 7 days following the procedure

#### Post-procedure dressing care

• After 24 hours you may remove any bandages and shower.

## Post-procedure discomfort or concerns

- For discomfort, you may take any NSAID (ex. **Motrin 600mg**) every 8 hours. Continue Motrin for 3-4 days post-procedure as needed to decrease the inflammation. If you are unable to take Motrin, you can take Tylenol.
- You may notice several "lumps" after the procedure. These may or may not be tender, and may appear red or pink.
  - Do not be concerned as this occurs in 1/3 of patients after the procedure. They are segments of residual veins with superficial clots
    that are not dangerous and will resolve. Apply warm compresses to these areas several times a day, and start taking motrin twice a
    day for 3-4 days.
- If you experience oozing/bleeding through your bandages, apply pressure with gauze using 2 fingers and lie
  down with your legs elevated. This will stop the bleeding almost always. If there continues to be bleeding call
  907-339-9455

\*For significant bleeding, fevers, signs of infection or any concerns contact your physician or go to the nearest emergency room.

### Post-procedure travel

 Wait 1 week before flying or travelling long distances (> 2 hours), unless cleared to travel sooner by your physician.

## Follow-up after your procedure

- 3-7 days you will have a scheduled ultrasound imaging visit only
- 10-14 days initial post op visit with a provider

\*If any questions, please feel free to contact our office at 907-222-4624